

**How to cite this article:/Cómo citar este artículo**

· Villalba-Moreno A, Alfaro-Lara ER, Sánchez-Fidalgo S, Nieto-Martín MD,  
· Santos-Ramos B. Development of the Anticholinergic Burden Calculator  
· Web Tool. Farm Hosp. 2017;41(5):647-648.



## CARTAS AL DIRECTOR

**Development of the Anticholinergic Burden Calculator Web Tool****Desarrollo de la herramienta web *Anticholinergic Burden Calculator***

Ángela Villalba-Moreno<sup>1</sup>, Eva Rocío Alfaro-Lara<sup>2</sup>, Susana Sánchez-Fidalgo<sup>3\*</sup>, María Dolores Nieto-Martín<sup>4</sup>, Bernardo Santos-Ramos<sup>1</sup>

<sup>1</sup>Pharmacy Service, University Hospital Virgen del Rocío, Seville. <sup>2</sup>Andalusian Agency for Health Technology Assessment, Seville. <sup>3</sup>Pharmacy Service, University Hospital Valme, Seville. <sup>4</sup>Internal Medicine Service. University Hospital Virgen del Rocío, Seville. Spain.

**Autor para correspondencia**

Correo electrónico:

susana.sanchez.exts@juntadeandalucia.es  
(Susana Sánchez-Fidalgo)

· Recibido el 24 de marzo de 2017;  
· aceptado el 28 de abril de 2017.  
· DOI: 10.7399/fh.10799

**Sra. Directora:**

The anticholinergic burden is defined as the cumulative effect of concomitantly taking multiple drugs with anticholinergic properties. It estimates the risk of suffering anticholinergic adverse effects. Anticholinergic scales are lists that rank the anticholinergic potential of drugs into categories. A patient's burden may be calculated from the sum of the score of each drug included in the scale. This assessment provides clinicians the opportunity to reconsider the pharmacotherapy in order to optimize treatment<sup>1</sup>.

Our research group has designed and developed a tool which calculates the anticholinergic burden based on 10 anticholinergic scales which have been firstly identified in a review<sup>2</sup> with the aim to be used in clinical setting. The tool was named *Anticholinergic Burden Calculator (ABC)*. It is a web portal software program ([www.anticholinergicscales.es/](http://www.anticholinergicscales.es/)). (An app for smart phones is now in progress).

Anticholinergic risk scores are defined into three groups, low, medium and high, according to risk categorization made by the authors of each scale. The user can view a report that contains three tables and one graphic that describe the results (Fig. 1). Drug Burden Index scale is separately described because it is the only one based on a simple mathematical formula that takes into account the prescribed dose and the minimum effective dose<sup>3</sup>.

The *Anticholinergic Burden Calculator (ABC)* has been validated for accuracy. Medication data including drug name and dose were collected from 35 patients with multimorbidity included in the IMPACTO project

involving a total of 310 drugs<sup>4</sup>. The anticholinergic burden was calculated twice, manually and with *Anticholinergic Burden Calculator (ABC)*. The results showed a 100% agreement.

The hand made anticholinergic burden calculation may be a tedious process. The tool facilitates the calculation of the total burden using 10 scales in a single step. Moreover, the professionals obtain very valuable information about treatment identifying which patients are at risk. On the other hand, they know drugs associated with increased risk and their contribution to the total load (the higher score, the greater risk). Thus, it facilitates decision-making on patient pharmacotherapy, reconsidering the treatment to reduce the load, for example, by deprescription, decreasing doses or substituting one drug for another without anticholinergic properties.

Our research group is currently working to improve the functionality of the web. Specific recommendations to help clinicians to make better decisions are in progress.

**Funding**

This study has been supported by grants from Consejería de Salud, Junta de Andalucía (PI-0195-2016).

**Conflicts of interests**

The authors concerned declare that they do not have any conflicts of interest.



Los artículos publicados en esta revista se distribuyen con la licencia  
Articles published in this journal are licensed with a  
Creative Commons Attribution 4.0  
<https://creativecommons.org/licenses/by-nc-nd/4.0/>  
La revista Farmacia no cobra tasas por el envío de trabajos,  
ni tampoco por la publicación de sus artículos.

### Anticholinergic burden results

The results of anticholinergic risk (low / medium/ high) obtained with each scale are linked to the risk categorization made by the authors or developers of each one of them

Scale	Result	Risk
ACB	0	WITHOUT RISK
ARS	0	WITHOUT RISK
Chew	0	WITHOUT RISK
ADS	2	WITHOUT RISK
AAS	0	WITHOUT RISK
ALS	0	WITHOUT RISK
CrAS	0	WITHOUT RISK
Duran	0	WITHOUT RISK
ABC	0	WITHOUT RISK

Medication	Scales									
	ACB	ARS	Chew	ADS	AAS	ALS	CrAS	Duran	ABC	
ALFUZOSIN 10 mg	0	0	0	0	0	0	0	0	0	
CORTISONE	0	0	0	1	0	0	0	0	0	
LORAZEPAM 5 MG	0	0	0	1	0	0	0	0	0	

**DBI Results** (Note: This scale, unlike the above, considers drug dose prescribed in the calculation)

Medication	DBI
ALFUZOSIN 10 mg	0.80
CORTISONE	0.00
LORAZEPAM 5 mg	0.83
<b>Result</b>	<b>HIGH RISK</b> 1.63




Figure 1. Tables and graphic according to risk score.

### References

- Durán CE, Azermi M, Vander Stichele RH. Systematic review of anticholinergic risk scales in older adults. *Eur J Clin Pharm.* 2013;69:1485-96.
- Villalba-Moreno AM, Alfaro-Lara ER, Pérez-Guerrero C, Nieto-Martín MD, Santos-Ramos B. Systematic review on the use of anticholinergic scales in poly pathological patients. *Arch Gerontol Geriatr.* 2016;62:1-8.
- Hilmer SN, Mager DE, Simonsick EM, Cao Y, Ling SM, Windham BG. A Drug Burden Index to define the functional burden older people. *Arch Intern Med.* 2007;167:781-7.
- Implantación del Plan de Asistencia Continuada a Pacientes Patológicos. Impacto sobre la Evolución Natural de la Enfermedad, el Deterioro Funcional y la Calidad de Vida Proyecto IMPACTO [Monografía en Internet]. [Consultado 20/03/2017]. Disponible en: <https://www.fesemi.org/grupos/edad-avanzada/proyectos/proyecto-impacto>