

How to cite this article:

· Ginestal-López RC. Immunotherapy for neurological diseases, present and future. Farm Hosp. 2018;42(6):251-260.



REVIEW

Bilingual edition english/spanish

Immunotherapy for neurological diseases, present and future

Inmunoterapia en enfermedades neurológicas, presente y futuro

Ricardo Constantino Ginestal-López

Neurology Department, Hospital Universitario Fundación Jiménez Díaz UTE, Madrid, Spain.

Author of correspondence

Ricardo C. Ginestal López
Servicio de Neurología, Hospital Universitario
Fundación Jiménez Díaz UTE.
Avenida de los Reyes Católicos 2, Madrid,
28040. Spain.

Email:
rcginestal@yahoo.es

· Recibido el 5 de abril de 2018;
· aceptado el 30 de julio de 2018.

· DOI: 10.7399/fh.11031

Abstract

Objective: The objective of this work is to summarize the immunological treatment of neurological diseases, describing the present situation and the challenges and opportunities the future will present.

Method: After topographically classifying the autoimmune neurological pathologies, a bibliographic analysis is made to present the most relevant ones regarding the available immunotherapeutic options. Likewise, new neurological entities that will be future candidates for immunotherapy are discussed.

Results: There is a large number of neurological diseases with an autoimmune basis, even though their pathophysiology is, sometimes, only partially understood. Only a few randomized controlled clinical trials support the evidence of the immunotherapies with which we treat some of these diseases. This situation is rapidly changing among entities like multiple sclerosis where level 1 of evidence clinical studies are today's standard. Alzheimer's disease and migraine are two of the most prevalent conditions that are being incorporated to the group of diseases candidates for immunotherapy.

Conclusions: Taking into account the rapidly growing number of immunological therapies and of neurological diseases potentially receiving them, an adequate evaluation of the impact these treatments will have on social and healthcare system grounds is necessary to reach compromises and consensus among all the professionals involved in the management of these pathologies.

Resumen

Objetivo: El objetivo del presente trabajo es resumir el tratamiento inmunológico de las enfermedades neurológicas, describiendo la situación actual y los retos y oportunidades que se presentan en un futuro próximo.

Método: Se realiza un análisis bibliográfico para, tras clasificar topográficamente las patologías neurológicas autoinmunes, presentar las más relevantes según las opciones inmunoterapéuticas disponibles. Asimismo, se exponen otras enfermedades neurológicas que serán nuevas candidatas a terapia inmunológica en el futuro.

Resultados: Existen múltiples patologías neurológicas con base autoinmune, aunque su fisiopatología, a veces, solo sea parcialmente conocida. Sin embargo, pocos son los ensayos clínicos aleatorizados y controlados que soportan la evidencia de los tratamientos inmunológicos con los que las tratamos. Esta situación está cambiando rápidamente en enfermedades como la esclerosis múltiple, donde ensayos clínicos con un nivel de evidencia grado 1 son la norma. La enfermedad de Alzheimer y la migraña son algunas de las más prevalentes que se están incorporando al grupo de candidatas a inmunoterapia.

Conclusiones: Con un número rápidamente creciente de terapias inmunológicas y de enfermedades neurológicas potencialmente tratables por esta vía, será necesaria una adecuada evaluación del impacto socio-sanitario que van a conllevar para llegar a compromisos y consensos por parte de todos los actores implicados en su manejo.

KEYWORDS

Immunotherapy; Neurologic disorders; Autoimmune diseases; Multiple sclerosis; Neuromyelitis optica; Alzheimer disease; Migraine; Movement disorders.

PALABRAS CLAVE

Inmunoterapia; Patologías neurológicas; Enfermedades autoinmunes; Esclerosis múltiple; Neuromielitis óptica; Enfermedad de Alzheimer; Migraña; Trastornos del movimiento.



Los artículos publicados en esta revista se distribuyen con la licencia
Articles published in this journal are licensed with a
Creative Commons Attribution 4.0
<https://creativecommons.org/licenses/by-nc-nd/4.0/>
La revista Farmacia no cobra tasas por el envío de trabajos,
ni tampoco por la publicación de sus artículos.

Introduction

During the past decades there has been a significant increase in our knowledge about the physiopathology of neurological conditions. Within these, diseases with immunological origin are responsible for major morbidity and even mortality, affecting the central nervous system (CNS) (multiple sclerosis [MS], neuromyelitis optica [NMO], limbic encephalitis...) and/or the peripheral system (Guillain-Barré syndrome [GBS], myasthenia gravis...) (Table 1). Overall, all these nosological conditions cause a major impact on patients and those around them but, given that some of them have a high prevalence, there can also be an impact for the public health system.

The constant study of new treatment targets has allowed a more efficient treatment of neurological diseases with immunological origin, with medications more specific for each condition, such as the case of MS, but it has also led to widening the range of diseases that can be potentially treated in this way. Thus, a deeper knowledge of the physiopathological mechanisms of conditions so prevalent as Alzheimer's disease or migraine, so far not included in the group of "immune-mediated diseases", has shown the role of immunomodulation treatments for their management, not only as hypotheses for the future but also as a short-term reality.

This article is intended as a brief review of the current situation of immunotherapy in neurological conditions, and to present some of the new therapeutic targets that will be offered to us in the near future.

Multiple sclerosis

MS is an inflammatory and demyelinating condition of the CNS. Its prevalence in Spain is of 91.2/100,000¹, and there has been a confirmed increase in its prevalence during the past decades. It presents two evolution forms. The relapsing remitting form (RRMS) consists in the presence of relapses of focal and acute CNS inflammation which cause new symptoms, with or without irreversible accumulated sequelae (and without any clinical worsening when there are no relapses). The progressive form is defined by a worsening in the clinical status of the patient when there are no relapses. The treatment of the disease, besides symptomatic treatment and physiotherapy, is divided into the management of relapses and the use of "disease modifying drugs".

Relapses are treated with glucocorticoids at high doses in short courses, typically 1 daily gram of intravenous methylprednisolone during 3 to 5 days. A multicenter, randomized, double-blind controlled clinical trial confirmed the non-inferiority of oral methylprednisolone 1,000 mg/day/3 days

compared with intravenous methylprednisolone 1,000 mg/day/3 days²; therefore, oral treatment could be an alternative option to intravenous administration. In cases resistant to corticoid therapy, a plasmapheresis course will be effective in 72% of cases³.

Disease-modifying treatment is intended to improve the functional prognosis of patients at medium and long term. Until the launch of immunomodulatory treatments specific for MS management, treatment consisted in immunosuppressants used for other conditions. In fact, azathioprine and mitoxantrone have the approved indication for their use in this disease. When interferon-beta 1b appeared in the 90's decade, there was a revolution in MS treatment. Currently we have 17 different products for treatment of relapsing remitting MS, three for the treatment of secondary progressive forms, and one recently approved for the treatment of the primary progressive disease. All these have demonstrated their efficacy in randomized and controlled clinical trials, and this has led to the approval of the specific indication in their product specifications. Figure 1 shows the molecules used for treatment of MS, according to their order of launch. It is worth highlighting the increasing frequency with which new molecules for this indication are being approved by regulatory agencies.

Both interferons- β and glatiramer acetate are being used for the past two decades with moderate efficacy and a verified safety profile, making them adequate as first line treatment. There is only correct acceptance by patients, because the way of administration is subcutaneous or intramuscular. For this reason, there has been a search for formulations with lower frequency of administration, such as pegylated interferon- β or glatiramer acetate 40 mg. Widely used, this type of medications is not always effective or well tolerated in all patients; therefore, research has focused on finding medications which are more potent and ways of administration which are more convenient for patients.

Teriflunomide and dimethyl fumarate, which are oral medications, currently allow treating MS patients without using parenteral administration. Teriflunomide⁴ acts by a reversible inhibition of the dihydroorotate dehydrogenase enzyme, highly expressed in activated lymphocytes, causing a reduction in the proliferation of activated lymphocytes T and B. Teriflunomide has demonstrated efficacy in RRMS treatment both in controlled and randomized clinical trials vs. placebo and in the clinical trial vs. interferon β 1a 44 μ g three times per week. Its safety profile is favorable but there is potential liver toxicity, which requires strict monitoring. It cannot be used in pregnant women until two years after discontinuation, because there is high teratogenic risk; there is a procedure for fast elimination which allows to reduce the product concentration below the risk considered minimal for the fetus (<0.02 μ g/mL). Dimethyl fumarate⁵, with a mechanism of action still not completely understood, has demonstrated its efficacy in two clinical trials. Its gastrointestinal tolerability can be troublesome, unlike its other characteristic side effect: flushing. Some cases of progressive multifocal leukoencephalopathy (PML) have been published in association with this drug, which forces to conduct a periodical monitoring of the total lymphocyte count; it is recommended to interrupt the use of dimethyl fumarate if this count goes below 500/ μ L persistently during 6 continuous months. In terms of finding more effective drugs, medications such as alemtuzumab, fingolimod and natalizumab are those associated with higher efficacy for preventing relapses; natalizumab seems to be associated to a higher extent with a lower progression of disability⁶. Natalizumab is a recombinant humanized monoclonal antibody which binds with integrin alpha-4 beta-1 and blocks the interaction with the vascular cell adhesion molecule 1 (VCAM-1), thus preventing the migration to the CNS of mononuclear leukocytes through the endothelium of the blood-brain barrier. Its use has been associated with the development of PML in patients with positive test results for anti-JC virus antibodies, and this risk will be higher with a longer time of treatment, and also if the patient was treated with immunosuppressants before natalizumab. This fact is a major limitation for the use of this medication. Alemtuzumab has been evaluated in three clinical trials vs. an active comparator: interferon β 1a 44 μ g three times per week. A Cochrane evaluation reached the conclusion that alemtuzumab reduces the percentage of patients who suffer relapses, disability progression and development of new lesions, as seen in magnetic resonance imaging throughout 24 to 36 months, compared with interferon β 1. Alemtuzumab has not been associated with PML development; but there have been potentially severe reactions to the intravenous infusion, infec-

Table 1. Preferential location of the autoimmune neurological damage and associated conditions.

Location	Condition
The whole central nervous system	Acute disseminated encephalomyelitis Multiple sclerosis
Cerebral cortex	Limbic encephalitis Autoimmune epilepsy Autoimmune dementia
Midbrain (Diencephalon)	Hypothalamic dysfunction
Basal ganglia	Chorea/Dystonia/Dyskinesia
Cerebellum	Cerebellar ataxia
Brain stem	Brain stem encephalitis Stiff-person syndrome
Cranial nerves	Isolated or multiple neuropathies
Spinal cord	Myelopathy and myoclonus
Peripheral nerves and nodes	Sensory neuronopathy and motor and sensory neuropathy
Motor plaque	Myasthenia gravis Eaton-Lambert syndrome
Muscle	Polymyositis/Dermatomyositis/Necrotizing Myopathy/Inclusion-body myositis
Autonomic nervous system	Dysautonomia and bowel movement alterations




Figure 1. Molecules approved for Multiple Sclerosis treatment in chronological order of availability for clinical practice.

tions, and autoimmune events which require strict monitoring. Fingolimod is a sphingosine-1-phosphate receptor modulator with oral administration, which has demonstrated its efficacy and safety for RRMS treatment in three Phase III clinical trials. It is more effective than weekly interferon β 30 μ g in the reduction of relapse parameters and magnetic resonance imaging. Given that fingolimod also interacts with different subtypes of sphingosine-1-phosphate receptors (S1PR1, S1PR2, S1PR3, S1PR4, S1PR5), there is a risk of bradycardia and QT-interval prolongation, which requires patient monitoring during the first dose. An increase in blood pressure has been described, as well as macular edema, liver toxicity and some cases of PML, which require a careful monitoring of patients⁸. With the aim to improve the safety profile of sphingosine-1-phosphate receptor modulators, there are various molecules under research with higher selectivity for the S1PR1 receptor, which is the cause of the effect on lymphocytes, and has no effects on other organs or systems. The agents currently under development are: siponimod, ponesimod, ozanimod, ceralifimod, GSK2018682 and MT-1303. Siponimod stands out among these, because it has demonstrated, in a Phase III clinical trial on the secondary progressive form of MS vs. placebo, its ability to reduce the confirmed progression disability at 3 and 6 months by 21% and 26%, respectively⁹. These results could lead siponimod to become, alongside interferon β 1a 44 μ g, interferon β 1b and mitoxantrone, the fourth medication with specific indication for the treatment of secondary progressive MS.

In this overview of the near future of MS treatment, we cannot leave out two medications recently approved by the European Medicines Agency: ocrelizumab and oral cladribine. Ocrelizumab is a humanized anti-CD20 monoclonal antibody that has demonstrated high efficacy in RRMS treatment, based in two randomized and controlled clinical trials. It has also confirmed, and this is a milestone in MS treatment, its ability to delay the accumulated disability in patients with primary progressive MS¹⁰. Thus, it has the indication for RRMS but also for the treatment of those adult patients with early primary progressive MS who present inflammatory activity in imaging tests. Other anti-CD20 antibodies have already been used off-label for RRMS treatment (rituximab), or are currently in the stage of clinical development (ofatumumab). Oral cladribine¹¹ causes gradual lymphocyte depletion over the weeks, not associated with cell lysis, with higher impact on B cells than on T cells, and with reconstitution of the count of said cell lines throughout the months. In this way, when administered in two bimonthly cycles separated by one year, it acts as an inductor drug, not causing the prolonged immunosuppression of other previously mentioned medications which require uninterrupted treatment. The efficacy of cladribine has been demonstrated in two randomized and controlled Phase III clinical trials. Its main side effect is lymphopenia, associated with its mechanism of action; but it does not seem to be associated with an increase in neoplasia or infections vs. the control, except for the case of herpes zoster infections. Cla-

drine is indicated for adult patients with recurrent MS with high clinical or radiological activity.

The way of administration, potential effects, and recommended monitoring for the medications mentioned in this review appear summarized in table 2.

Neuromyelitis optica

NMO is an inflammatory demyelinating condition, anti-aquaporin 4 antibody-mediated (NMO-IgG). It affects specifically the spinal cord and the optic nerves. Its prevalence in Spain is of 1.5/100,000 inhabitants, and therefore less frequent than MS, but potentially more severe in the majority of cases.

Treatment of inflammatory relapses is conducted with 1 gram of intravenous methylprednisolone per day during 3 to 5 days, though its evidence comes from studies with MS or optic neuritis patients. Those patients who show no improvement with the previous regimen are treated with plasmapheresis¹² or intravenous human immunoglobulin¹³.

Regarding maintenance treatment, intended to prevent new relapses and accumulated disability, treatment will be typically initiated with azathioprine or mycophenolate mofetil, while the patient receives treatment with IV methylprednisolone, due to the time that these medications will take to start acting. Another first line option is rituximab¹⁴. Methotrexate would be reserved for those patients who don't tolerate the previous treatments, or those for whom these are not effective¹⁵. Other potential treatments, but more dubious in terms of efficacy or toxicity, are tacrolimus, cyclosporine, mitoxantrone, and cyclophosphamide¹⁶. It is worth pointing out here that some disease-modifying drugs for MS, such as interferon β ¹⁷, natalizumab¹⁸, and fingolimod¹⁹, will lead to a worsening in the evolution of NMO; therefore, it is essential to conduct an adequate differential diagnosis between both nosological entities.

In terms of new treatment options, various monoclonal antibodies are being evaluated for their use in NMO. Tocilizumab is a recombinant humanized anti-IL-6, which causes deletion of plasmablasts (which are CD20, and therefore are not affected by rituximab); it has shown efficacy in some isolated cases²⁰⁻²⁴ and in a Phase IV study²⁵. On the other hand, eculizumab inhibits the complement pathway, preventing the cleavage of C5 into C5a and C5b, and therefore preventing the formation of membrane attack complex (C5b-C9). This molecule has shown potential for NMO treatment in a pilot open study with 14 patients²⁶.

Other potentially valid treatment options, still pending adequate assessment, would be: aquaporin 4 antibody (a humanized NMO-IgG monoclonal antibody with high affinity which might prevent pathogenic aquaporin-4 from binding to the Fc fragment), alemtuzumab (anti-CD52 previously

Table 2. Immunomodulatory treatments: way of administration, potential risks, and recommended monitoring.

Treatment	Way of administration	Potential risks	Monitoring
Glucocorticoids	IV in the acute stage Oral for maintenance	Hyperglycemia, hypertension, sleep and behaviour alterations, adrenal suppression, osteoporosis, osteonecrosis, myopathy, glaucoma, cataracts, GI alterations	Blood pressure, glycemic profile, ophthalmological evaluation, bone densitometry
Immunoglobulins	IV SC	Hypersensitivity alterations, thromboembolic events, renal failure, aseptic meningitis, hemolytic anemia, neutropenia	Vital signs during infusion, BUN and creatinine
Plasmapheresis	IV	Central line, hypocalcemia, arterial hypotension, arrhythmia, coagulopathy	Blood test, electrolytes, immunoglobulin levels, coagulation
Azathioprine	Oral	Liver toxicity, cytopenias, GI toxicity, alopecia, photosensitivity, risk of lymphoma	Blood test, liver profile, 5-thiopurine methyltransferase testing before treatment
Cyclophosphamide	IV, oral	Nausea/vomiting, alopecia, mucositis, cystitis, teratogenicity, egg/spermatozoid preservation due to lack of fertility	Blood test, renal function, urinetest
Methotrexate	Oral	Nausea, diarrhea, mucositis, cytopenias, liver toxicity, pneumonitis, photosensitivity, teratogenicity	Blood test, liver profile, chest X-ray before treatment
Mitoxantrone	IV	Injection-site necrosis, arrhythmias, cardiopathy, cytopenias, liver toxicity, egg/spermatozoid preservation due to lack of fertility, leukemia	Not exceeding the maximum dose of 140 mg/m ² , blood test, liver profile, ECG, echocardiogram
Mycophenolate mofetil	Oral	Nausea, diarrhea, abdominal pain, liver toxicity, cytopenias, hypertension, nephrotoxicity, coughing, dyspnea, headache, tremor, lymphoma and other neoplasia, teratogenicity	Blood test, liver profile, renal function
Tacrolimus	Oral	Headache, nephrotoxicity, hypertension, tremor, diarrhea, nausea/vomiting, insomnia, anorexia, distal paresthesias	Blood pressure, renal function
Interferon-β	SC, IM	Flu-like syndrome, injection-site reactions, uremic hemolytic syndrome, dysthyroidism, liver toxicity, depression	Blood test, renal function, liver profile, thyroid function
Glatiramer acetate	SC	Injection-site reactions, lipoatrophy, post-injection idiosyncratic reaction, lymphadenopathy	Not required
Fingolimod	Oral	PML, bradycardia at first dose, hypertension, lymphopenia, liver toxicity	Previous encephalic MRI, cardiac monitoring at first dose, blood test, liver profile, blood pressure
Teriflunomide	Oral	Liver toxicity, cytopenias, hypertension, lack of GI tolerability, alopecia, peripheral neuropathy, teratogenic	Blood pressure, blood test, liver profile
Dimethyl fumarate	Oral	Lack of GI tolerability, flushing, PML, cytopenia, liver toxicity	Previous encephalic MRI, blood test, liver profile
Natalizumab	IV	PML, hypersensitivity alterations at infusion	Previous encephalic MRI, and periodical follow-up MRI, blood test, liver profile, anti-JCV testing every 6 months in seronegative patients
Alemtuzumab	IV	Hypersensitivity alterations at infusion, thrombopenia, dysthyroidism, glomerulonephritis	Blood test, liver profile, renal function, urine test, vital signs during infusion
Rituximab	IV	Hypersensitivity alterations, hypogammaglobulemia, PML, infusion reactions, edema, fever, headache	Vital signs during infusion, blood test, CD19/CD20 count, IgG/IgM levels
Ocrelizumab	IV	Hypersensitivity alterations at infusion, infections, neutropenia, reduction in IgM and IgG, coughing, ENT mucosity, cancer (breast)	Blood test, liver profile
Cladribine	Oral	Cytopenias (lymphopenia), liver toxicity, infections (herpes zoster)	Blood test, liver profile, watching for neoplasia
Tocilizumab	IV	Hypersensitivity alterations, GI perforation, liver toxicity, neutropenia, thrombopenia, TB reactivation	Blood test, liver profile
Eculizumab	IV	Hypersensitivity alterations, hypertension, anemia, meningococcal infection	Blood test, liver and renal profile
Infliximab	IV	Hypersensitivity alterations, liver toxicity, demyelination, Neuritis optica, TB reactivation	Vital signs during infusion, blood test, liver profile

ABUN: Blood urea nitrogen test; MRI: Magnetic resonance imaging; GI: Gastrointestinal; TB: Tuberculosis; PML: Progressive multifocal leukoencephalopathy; ENT: Ear, nose and throat; ECG: Electrocardiogram; IgG: Immunoglobulin G; IgM: Immunoglobulin M; Anti-JCV: JC virus antibody test.

described in the treatment for MS), and infliximab (chimeric monoclonal antibody anti-TNF α)²⁷.

Migraine

Migraine is a highly prevalent neurological disease; 14.7% of the world population suffers it. It is the third human condition more frequent after tooth cavities and tension headache²⁸. Its treatment is based on stopping the acute pain attack (with non-steroid anti-inflammatories and triptans, mainly); but also on medications to prevent new episodes, in those patients where the frequency and intensity of pain will justify their use. Currently, anti-hypertension drugs are used (betablockers), as well as antidepressants (tricyclics) and antiepileptic agents (topiramate, valproic acid), among others²⁹. In some cases, botulinum toxin has also been used³⁰⁻³².

In the past decade, blocking the calcitonin gene-related peptide (CGRP) has also been put forward as a new treatment target for preventing migraine episodes. This is due to the finding that CGRP levels increased during migraine episodes, and were reduced after using triptans³³. Moreover, the intravenous administration of CGRP caused migraine episodes in migraine patients³⁴. Two types of drugs have been designed in order to modify this CGRP design: on one hand, the CGRP-receptor antagonists, the "gepant" class, and on the other hand, monoclonal antibodies targeted to CGRP or its receptor. All these have demonstrated efficacy in controlled and randomized clinical trials, with efficacy at least equal than that of current preventive treatments. Figure 2 summarizes the efficacy in clinical trials of monoclonal antibodies as prophylaxis for episodic and chronic migraine³⁵⁻⁴⁰.

In terms of safety, liver toxicity problems have been observed with some gepants; this does not seem to happen with monoclonal antibodies⁴¹. Regarding the latter, in the absence of long-term data, some doubts can arise in terms of the cardiovascular system (hypertension, ischemic events), pituitary function, the GI system (constipation/diarrhea, ulcers, irritable bowel), and skin (erythema, inflammation, interference with wound healing)⁴¹. The potential presence of neutralizing antibodies which limit the effect of monoclonal antibodies over time, and the cost of treatment, can also be factors to take into account regarding their use in a disease with an extraordinary prevalence.

Movement disorders

Movement disorders caused by an autoimmune physiopathology can appear isolatedly or within a wider encephalopathic process including epileptic manifestations and/or cognitive deterioration. Traditionally, movement disorders have been classified into "hyperkinetic" (myoclonus, chorea, tics, pseudoathetosis, dystonia and other phenomena), and "hypokinetic" (Parkinsonism, stiff-person syndrome, progressive encephalomyelitis with rigidity and myoclonus). These are conditions with low prevalence (e.g. stiff-person: 1/1,250,000). In their treatment, besides determining if there is a causal neoplasia and eliminating it, treatment consists in the use of intravenous methylprednisolone, intravenous human immunoglobulin, or plasmapheresis in the acute stage, and azathioprine and mycophenolate mofetil as maintenance therapy^{42,43}.

In this section, it is worth referring briefly to potential treatment targets in idiopathic Parkinson's disease, a condition with a major prevalence (1-2/1,000)⁴⁴. Physiopathological knowledge leads to consider that α -synuclein is a key molecule in neuronal death in Parkinson's disease, by aggregating in toxic forms, spreading into the extracellular space and "contaminating" the adjacent neurons, thus perpetuating the pathogenic process. Thus, active or passive immunotherapy strategies intended to reduce the level of α -synuclein toxic extracellular aggregation could reduce or prevent the disease progression^{45,46}.

Autoimmune epilepsy

Alongside epilepsy conditions associated with systemic autoimmune disorders, such as systemic lupus erythematosus, Hashimoto encephalopathy, sarcoidosis or celiac disease, there are antibody-mediated disorders which cause epileptic episodes as one of their main clinical manifestations. Said antibodies can be classified into those that bind intracellular antigens and those that bind to neuronal surface proteins. The first class includes the Hu, Ma2, CRMP5 and amphiphysin antibodies. Those conditions derived of intracellular antigens will present a worse response to immunotherapy⁴⁷. Overall, immune treatment consists in a first line with intravenous methylprednisolone 500-1,000 mg/day/5 days, while intravenous immunoglobulin or plasmapheresis are reserved for steroid-resistant patients⁴⁸. Second line




Figure 2. Efficacy of monoclonal antibodies in preventive treatment for migraine: reduction in migraine days per month (mean). All outcomes are statistically significant ($p < 0.05$). IV: intravenous administration; SC: subcutaneous administration.

treatment must be initiated within 2 weeks at most, if there has been no >50% reduction in episodes with first line agents^{48,49}. The following will then be used: cyclophosphamide⁴⁸, rituximab⁵⁰, cyclophosphamide + rituximab, mycophenolate mofetil, or azathioprine^{48,49}.

Dementia and autoimmune encephalopathies

The presentation form ranges from acute limbic encephalitis to subacute or chronic forms with a difficult differential diagnosis vs. primarily neurodegenerative conditions. Their etiology is primarily idiopathic-autoimmune or in the context of a paraneoplastic phenomenon. In terms of treatment, the main objective in paraneoplastic conditions is to remove the causal tumour completely, whenever possible. For any of both physiopathological mechanisms, however, acute treatment of neurological symptoms will be, as in other autoimmune conditions, intravenous methylprednisolone used at high doses, or intravenous human immunoglobulin. Improvement with acute treatment can justify the use of maintenance therapy with corticosteroids, azathioprine, mycophenolate mofetil, methotrexate or tacrolimus, among other agents. In some conditions, such as anti-N-methyl-d-aspartate receptor encephalitis, drugs like rituximab or cyclophosphamide can be considered second-line treatment when there is little or no response with previously mentioned agents^{51,52}. Acute disseminated encephalomyelitis, more frequent among pediatric patients, is a condition more typically monophasic, and therefore its management will usually be limited to acute treatment with intravenous methylprednisolone at high doses, intravenous human immunoglobulin, or plasmapheresis⁵³.

Alzheimer's disease

The prevalence of Alzheimer's disease increases with age. In an estimation conducted for 2017 in the United States of America, 4% of people <65-year-old present the disease, as well as 16% of persons between 65 and 74-year-old, 44% between 75 and 84-year-old, and 38% of people >85-year-old. It is expected that, at world level, the number of Alzheimer cases will triple by 2050⁵⁴. In an ageing society like ours in Spain, Alzheimer's disease is already representing a major public health problem, and this will get even worse in the future.

The treatment for Alzheimer's disease is based on those items considered key in its physiopathology, still only partly known (Table 3). A key event in this condition is the loss of cholinergic neurons, which has led to the prescription of cholinesterase inhibitors such as donepezil or rivastigmine. Another relevant element is the increase in glutamatergic activity, which has strengthened the indication of anti-N-methyl-D-aspartate receptor blockers such as memantine. Both therapeutic classes are typically used in current clinical practice. There are other three physiopathological mechanisms that can open new therapeutic targets, this time through the use of immunotherapies. The build-up of neuritic plaques (β -amyloid), neurofibrillary tangles (τ protein), and the local inflammation caused by microglia, are potential treatment objectives that are being evaluated in recent years⁵⁵.

β -amyloid plaque is formed in two steps from the amyloid precursor protein (APP), through the β -secretase and γ -secretase enzyme complexes. It is thought that β -amyloid deposit could play a major role in the development of Alzheimer's disease⁵⁶. Keeping this hypothesis in mind, there has been a development, on one hand, of therapies able to reduce the activity of the β -secretase and γ -secretase complexes, with the objective of producing less β -amyloid; and, on the other hand, immunological treatments that can increase the elimination of the pathogenic β -amyloid already formed.

Thus, two strategies have been developed: active immunotherapy and passive immunotherapy. One of the first clinical trials on active immunotherapy was conducted with AN1792, a synthetic amyloid peptide (A β 42) which induces the production of antibodies against the β -amyloid⁵⁷. The Phase II clinical trial was stopped due to the development of meningoencephalitis in 6% of the study subjects. Besides, in those who did not develop meningoencephalitis, there was no demonstrated delay in the evolution of cognitive deterioration, despite a clear reduction in the senile plaque deposits. In view of these results, it was argued that the study included patients with moderate/severe Alzheimer's disease; maybe treatment with drugs promoting the elimination of senile plaques could be indicated in earlier stages of the disease. With this recommendation, other molecules promoting active immunity to eliminate the amyloid plaques are being evaluated: among them, agent CAD106, safe and well tolerated, without any meningoencephalitis cases reported to date, is still under study⁵⁸. Regarding passive immunotherapy, various molecules have been developed and are currently under evaluation. Bapineuzumab, targeting the N-terminal end of β -amyloid, has been discontinued due to its low efficacy and negative safety profile at Phase III^{59,60}. On the other hand, solanezumab, which targets the monomer soluble β -amyloid, has presented an adequate safety profile, and it is being evaluated in patients with very early forms of Alzheimer's disease. It has demonstrated an increase in β -amyloid-42 soluble protein in the cerebrospinal fluid of patients, with dose-dependent effect, which would support its mechanism of action of β -amyloid deposit elimination. However, it has not demonstrated clinical efficacy in mild Alzheimer's disease⁶¹. Aducanumab is also on Phase III clinical trials, used for mild Alzheimer's disease, but also for the mild cognitive impairment stage. Finally, gantenerumab, which interacts with β -amyloid fibrils to recruit microglia, activate phagocytosis and degrade neuritic plaques, is currently in early stages of clinical development; it has shown an adequate safety profile, and is pending efficacy outcomes in mild Alzheimer's disease and mild cognitive impairment⁶².

Similarly as with neuritic plaques, the presence of neurofibrillary tangles formed by hyperphosphorylated τ protein seems to be one of the key elements responsible for pathogenesis in Alzheimer's disease. Aggregated τ protein is cytotoxic; therefore, preventing its production or favoring its elimination could have a clinical effect on patients. As it has been mentioned for β -amyloid protein, there are molecules which could lead to the elimination of pathogenic τ protein, through active immunotherapy. AADvac-1, an active vaccine with a natural truncated form of τ , is currently on Phase II

Table 3. Key physiopathological elements in Alzheimer's disease and treatment strategies targeting them

		Key physiopathological elements				
		Loss of cholinergic neurons	Increase in the glutamate activity	Build-up of neuritic plaques (β -amyloid deposit)	Neurofibrillary tangles (cytotoxic aggregated τ protein)	Increase in neuroinflammation (microglia activation)
Treatment strategies	Acetylcholinesterase Inhibitors			β -secretase inhibitors	\uparrow τ -protein stabilization	
	Serotonergic transmission modulators			γ -secretase inhibitors	τ -protein aggregation inhibitors	
	Histaminergic transmission modulators		N-methyl-d-aspartate blockers			\downarrow microglia activation
	\uparrow response to acetylcholine			\uparrow β -amyloid clearance, IMMUNOTHERAPY (active and passive)	\uparrow τ -protein clearance, IMMUNOTHERAPY (active and passive)	IMMUNOTHERAPY

after having shown a good safety profile. C2N8E12 is a humanized anti- τ antibody currently on Phase II study, and safety and efficacy outcomes are expected for 2020⁶³.

Besides neuritic plaques (β -amyloid) and neurofibrillary tangles (τ), neuroinflammation is one of the key elements in the physiopathology of Alzheimer's disease. Evident astrogliosis has been observed, among other signs of inflammation, around amyloid plaques, and various studies suggest a relationship between microglia activation, neuritic plaque formation, and the clinical progression of the disease. For this reason, a therapy targeted to inhibit the activation of microglia could be useful. However, negative results with tramiprosate (lack of efficacy at Phase III), ibuprofen and r-flurbiprofen, have reduced the expectations for this therapeutic group / mechanism of action. CHF 5074 is still on Phase II clinical trials: a microglia modulator for patients with mild cognitive impairment⁶³.

Finally, it is worth pointing out that intravenous human immunoglobulin seems to be effective for maintaining the cognitive ability in patients with mild to moderate Alzheimer's disease in Phase III clinical trials⁶⁴.

Peripheral nervous system

- GBS: An acute sensory-motor polyradiculopathy, demyelinating, axonal or mixed, with inflammatory origin. The typical symptoms are paresthesia, pain and loss of strength, though there are different clinical forms such as Miller Fisher syndrome, a sensory ataxia condition with involvement in the brain stem and oculomotor nerves. Its incidence is of 0.4 to 3.25 patients per 100,000 inhabitants and year⁶⁵. GBS is treated with plasmapheresis; its efficacy was confirmed in Cochrane's 2012 review⁶⁶. An alternative option is using intravenous human immunoglobulin at 0.4 g/kg/day during 5 days. No placebo-controlled studies have been conducted, but a Cochrane review from 2014 confirmed its efficacy, comparable to plasmapheresis, after a systematic evaluation of 5 clinical trials⁶⁷. Besides, two clinical trials confirmed similar efficacy but fewer side effects of immunoglobulin vs. plasmapheresis^{68,69}. A complete review on this matter can be consulted in the article by Wijdicks and Klein⁷⁰.
- Chronic Inflammatory Demyelinating Polyneuropathy (CIDP): Considered the chronic presentation (> 8 weeks of duration) of the demyelinating form of GBS, its treatment also consists in the use of plasmapheresis or immunoglobulin. However, while glucocorticoids are considered ineffective in GMS, these are useful both in the acute stage and the maintenance treatment of CIDP⁷¹. Rituximab has shown efficacy in studies with small patient cohorts with this condition^{72,73}. Eculizumab could be an option still not tested in CIDP. Fingolimod⁷⁴ and alemtuzumab⁷⁵ could play a role in the treatment of this disease, pending confirmation in randomized and controlled clinical trials.
- Multifocal motor neuropathy (MMN) is an autoimmune disorder with low prevalence (0.6-2 patients per 100,000 inhabitants)⁷⁶. It causes a slowly progressive loss of strength, asymmetrical and mainly distal. MMN is mediated by antiganglioside antibodies, and it can be treated with human intravenous or subcutaneous immunoglobulin, thus improving its symptoms and preventing their progression. Four clinical trials⁷⁷⁻⁸⁰ have demonstrated that 78% of patients treated with intravenous immunoglobulin improved significantly their motor ability vs. 4% treated with placebo. Even though the meta-analysis of said studies did not show significant differences in the improvement of disability⁸¹, the treatment guidelines by the European Federation of Neurological Societies (EFNS) recommend using 2 g/kg of intravenous human immunoglobulin for first line treatment of MMN; this dose must be administered over 2 to 5 days. These guidelines also state that the maintenance dose of intravenous human immunoglobulin to be administered after an initial improvement with the first cycle should be 1 g/kg every 2 to 4 weeks, or 2 g/kg every 1-2 months⁸². It is worth noting that subcutaneous immunoglobulin has demonstrated similar efficacy to the intravenous formulation in MMN treatment, both for early stages⁸³ and as maintenance⁸⁴. For an extensive review about this, it is recommended to read the work by Kumar *et al.*⁸⁵.
- Amyotrophic lateral sclerosis (ALS) is a degenerative motor neuron disease. Physiopathological mechanisms of immunological substrate have recently been mentioned, thus opening a potential pathway for immunotherapy treatment. A review on this topic⁸⁶ describes the use of treatments for Rheumatoid Arthritis: anakinra (a recombinant analog of the interleukin-1 receptor antagonist), with negative outcomes; masitinib (a tyrosine-kinase inhibitor), with an on-going Phase III clinical trial, and tocilizumab, also under clinical trial, in this case Phase II. Treatments for MS have also been used, such as glatiramer acetate (negative outcomes) or fingolimod (efficacy not demonstrated). The lack of efficacy outcomes obtained with intravenous immunoglobulin, celecoxib, oza- nezumab, NPO01 (taurine), thalidomide, granulocyte stimulating growth factor, cyclosporine, or total lymphoid irradiation, have not prevented continuing with the line of research of immunological therapies for ALS. Other agents, such as ibudilast (a TL4 and phosphodiesterase 3 and 4 inhibitor), RNS60, or drugs used to prevent rejection in transplants (basaliximab + mycophenolate mofetil + tacrolimus + glucocorticoids) are currently under investigation.
- Myasthenia gravis I an antibody-mediated condition (anti-acetylcholine receptor -AChR- or anti-muscle specific kinase [MuSK]), which prevents an adequate transmission in the motor plaque. Its characteristic symptom is muscle fatigue. Initial treatment, besides acetylcholinesterase inhibitors, is based on the use of oral prednisone, intravenous immunoglobulin and/or plasmapheresis for disease relapses. In order to avoid the continuous use of glucocorticoids in patients with generalized disease, the following are used: azathioprine, mycophenolate mofetil, cyclosporine A, methotrexate, tacrolimus, or cyclophosphamide. For a complete review on the use of these medications for myasthenia gravis, the study by Lee and Jander⁸⁷ is recommended. Another treatment option is rituximab, an anti-CD20 chimeric monoclonal antibody, suggested for patients with moderate-severe forms of the disease who are refractory to other treatments, and for those who are anti-MuSK-positive^{88,89}. A meta-analysis evaluating 15 non-controlled clinical trials, with 168 patients included in total, on different treatment regimens with rituximab, seems to show its efficacy in the treatment of AChR-positive, MuSK-positive and AChR/MuSK-double negative myasthenia gravis⁹⁰. More recently, eculizumab has demonstrated efficacy in a Phase II clinical trial⁹¹ and in another in Phase III⁹²; for this reason, it could be considered initially as a treatment option for severe cases or those refractory to other treatment strategies⁹³.
- Autoimmune myopathies: The prevalence of polymyositis and dermatomyositis, the two most frequent autoimmune myopathies, is of 21.5/100,000 inhabitants⁹⁴. Corticosteroids are the first line treatment for both conditions, as well as for immune-mediated necrotizing myopathy⁹⁵⁻⁹⁷. In patients with severe symptoms (dysphagia or inability to walk), intravenous methylprednisolone is used at 1 g/day/3 days, followed by oral prednisone in a decreasing dose starting with 60 mg/day of prednisone. In moderate cases, it is possible to initiate oral treatment without the previous intravenous loading dose. In milder clinical presentations, it is possible to start at lower prednisone doses. Once muscular strength returns to normal, a progressive dose reduction will be implemented. In patients with severe disease, those with incomplete response to corticosteroids after 2 months of treatment, and those where the dose cannot be reduced below 10 mg/day, it is recommended to use azathioprine, methotrexate or mycophenolate mofetil. If there is failure or lack of tolerability to these second line medications, it could be possible to resort to rituximab, cyclosporine, cyclophosphamide, TNF α blockers⁹⁸. Intravenous human immunoglobulin is effective for dermatomyositis treatment⁹⁹ and possibly also for polymyositis. It can be used as second line for patients with severe symptoms, because its effect is faster than that of said treatment line. The typical dose is 2 g/kg, distributed between 3 to 5 days. Inclusion-body myositis does not respond to immunotherapy.

Discussion

Currently, immunological treatments allow us to treat a high number of neurological conditions in a more accurate and individualized way. The launch of products designed with specific treatment targets and evaluated in clinical trials with high level of evidence has become more frequent in clinical practice. Moreover, the range of conditions that can be potentially treated with immunotherapy is increasingly higher, and has started to include highly prevalent neurological conditions such as migraine and, possibly, Alzheimer's disease. The high impact that these therapies could entail for the

public health system will require compromises and consensus by all actors involved in their adequate use.

Funding

No funding.

Bibliography

- Otero-Romero S, Roura P, Solà J, Altimiras J, Sastre-Garriga J, Nos C, *et al.* Increase in the prevalence of multiple sclerosis over a 17-year period in Osona, Catalonia, Spain. *Mult Scler.* 2013;19(2):245-8.
- Le Page E, Veillard D, Laplaud DA, Hamonic S, Wardi R, Lebrun C, *et al.* Oral versus intravenous high-dose methylprednisolone for treatment of relapses in patients with multiple sclerosis (COPOUSEP): a randomised, controlled, double-blind, non-inferiority trial. *Lancet.* 2015;386(9997):974-81.
- Ehler J, Koball S, Sauer M, Mitzner S, Hickstein H, Benecke R, *et al.* Response to therapeutic plasma exchange as a rescue treatment in clinically isolated syndromes and acute worsening of multiple sclerosis: a retrospective analysis of 90 patients. *PLoS One.* 2015;10(8):e0134583.
- Miller AE. Oral teriflunomide in the treatment of relapsing forms of multiple sclerosis: clinical evidence and long-term experience. *Ther Adv Neurol Disord.* 2017;10(12):381-96.
- Linker RA, Haghighi A. Dimethyl fumarate in multiple sclerosis: latests developments, evidence and place in therapy. *Ther Adv Chronic Dis.* 2016;7(4):198-207.
- Tramacere I, Del Giovane C, Salanti G, D'Amico R, Filippini G. Immunomodulators and immunosuppressants for relapsing-remitting multiple sclerosis: a network meta-analysis. *Cochrane Database Syst Rev.* Sep 18;(9):CD011381.
- Zhang J, Shi S, Zhang Y, Luo J, Xia Y, Meng L, *et al.* Alemtuzumab versus interferon beta 1a for relapsing-remitting multiple sclerosis. *Cochrane Database Syst Rev.* Nov 27;(11):CD010968.
- Chaundry BZ, Cohen JA, Conway DS. Sphingosine 1-Phosphate receptor modulators for the treatment of multiple sclerosis. *Neurotherapeutics.* 2017;14(4):859-73.
- Kappos L, Bar-Or A, Cree BAC, Fox RJ, Giovannoni G, Gold R, *et al.* Siponimod versus placebo in secondary progressive multiple sclerosis (EXPAND): a double-blind, randomised, phase 3 study. *Lancet.* 2018;391(10127):1263-73.
- Gelfand JM, Cree BAC, Hauser SL. Ocrelizumab and other CD20+ B-cell-depleting therapies in multiple sclerosis. *Neurotherapeutics.* 2017;14(4):835-41.
- Giovannoni G. Cladribine to treat relapsing forms of multiple sclerosis. *Neurotherapeutics.* 2017;14(4):874-87.
- Weinshenker BG, O'Brien PC, Petterson TM, Noseworthy JH, Lucchinetti CF, Dodick DW, *et al.* A randomized trial of plasma exchange in acute central nervous system inflammatory demyelinating disease. *Ann Neurol.* 1999;46(6):878-86.
- Elsone L, Panicker J, Mutch K, Boggild M, Appleton R, Jacob A. Role of intravenous immunoglobulin in the treatment of acute relapses of neuromyelitis optica: experience in 10 patients. *Mult Scler.* 2014;20(4):501-4.
- Flanagan EP. Autoimmune myelopathies. *Handb Clin Neurol.* 2016;133:327-51.
- Kitley J, Elson L, George J, Waters P, Woodhall M, Vicent A, *et al.* Methotrexate is an alternative to azathioprine in neuromyelitis optica spectrum disorders with aquaporin-4 antibodies. *J Neurol Neurosurg Psychiatry.* 2013;84(8):918-21.
- Flanagan EP, Weinshenker BG. Neuromyelitis spectrum disorders. *Curr Neurol Neurosci Rep.* 2014;14(9):483.
- Palace J, Leite MI, Nairn A, Vincent A. Interferon beta treatment in neuromyelitis optica: increase in relapses and aquaporin 4 antibody titers. *Arch Neurol.* 2010;67(8):1016-7.
- Jacob A, Hutchinson M, Elson L, Kelly S, Ali R, Saulans I, *et al.* Does natalizumab therapy worsen neuromyelitis optica? *Neurology.* 2012;79(10):1065-6.
- Min JH, Kin BJ, Lee KH. Development of extensive brain lesions following fingolimod (FTY720) treatment in a patient with neuromyelitis optica spectrum disorder. *Mult Scler.* 2012;18(1):113-5.
- Kieseier BC, Stuve O, Dehmel T, Goebels N, Leussink VI, Mausberg AK, *et al.* Disease amelioration with tocilizumab in a treatment-resistant patient with neuromyelitis optica: implication for cellular immune responses. *JAMA Neurol.* 2013;70(3):390-3.
- Ayzenberg I, Kleiter I, Schröder A, Hellwig K, Chan A, Yamamura T, *et al.* Interleukin 6 receptor blockade in patients with neuromyelitis optica nonresponsive to anti-CD20 therapy. *JAMA Neurol.* 2013;70(3):394-7.
- Araki M, Aranami T, Matsuoka T, Nakamura M, Miyake S, Yamamura T. Clinical improvement in a patient with neuromyelitis optica following therapy with the anti-IL-6 receptor monoclonal antibody tocilizumab. *Mod Rheumatol.* 2013;23(4):827-31.
- Harmel J, Rigelstein M, Ingwersen J, Mathys C, Goebels N, Hartung HP, *et al.* Interferon-β-related tumefactive brain lesion in a Caucasian patient with neuromyelitis optica and clinical stabilization with tocilizumab. *BMC Neurol.* 2014;14:247.
- Komai T, Shoda H, Yamaguchi K, Sakurai K, Shibuya M, Kubo K, *et al.* Neuromyelitis optica spectrum disorder complicated with Sjögren syndrome successfully treated with tocilizumab: a case report. *Mod Rheumatol.* 2016;26(2):294-6.
- Ringelstein M, Ayzenberg I, Harmel J, Lauenstein AS, Lensch E, Stögbauer F, *et al.* Long-term therapy with interleukin 6 receptor blockade in highly active neuromyelitis optica spectrum disorder. *JAMA Neurol.* 2015;72(7):756-63.
- Pittock SJ, Lennon VA, McKeon A, Mandrekar J, Weinshenker BG, Lucchinetti GF, *et al.* Eculizumab in AQP4-IgG-positive relapsing neuromyelitis optica spectrum disorders: an open-label pilot study. *Lancet Neurol.* 2013;12(6):554-62.
- Lin J, Xue B, Li X, Xia J. Monoclonal antibody therapy for neuromyelitis optica spectrum disorder: current and future. *Int J Neurosci.* 2017;127(8):735-44.
- Steiner TJ, Stovner LJ, Birbeck GL. Migraine: the seventh disabling. *J Headache Pain.* 2013;14:1.
- Silberstein SD, Holland S, Freitag F, Dodick DW, Argoff C, Ashman E; Quality Standards Subcommittee of the American Academy of Neurology and the American Headache Society. Evidence-based guideline update: pharmacologic treatment for episodic migraine prevention in adults. *Neurology.* 2012;78(17):1337-45.
- Aurora SK, Dodick DW, Turkel CC, DeGryse RE, Silberstein SD, Lipton RB, *et al.*; PREEMPT 1 Chronic Migraine Study Group. OnabotulinumtoxinA for treatment of chronic migraine: Results from the double-blind, randomized, placebo-controlled phase of the preempt 1 trial. *Cephalalgia.* 2010;30(7):793-803.
- Diener HC, Dodick DW, Aurora SK, Turkel CC, DeGryse RE, Lipton RB, *et al.*; PREEMPT 1 Chronic Migraine Study Group. OnabotulinumtoxinA for treatment of chronic migraine: Results from the double-blind, randomized, placebo-controlled phase of the preempt 2 trial. *Cephalalgia.* 2010;30(7):804-4.
- Dodick DW, Turkel CC, DeGryse RE, Aurora SK, Silberstein SD, Lipton RB, *et al.*; PREEMPT 1 Chronic Migraine Study Group. OnabotulinumtoxinA for treatment of chronic migraine: Pooled results from the double-blind, randomized, placebo-controlled phases of the preempt clinical program. *Headache.* 2010;50(6):921-36.
- Goadsby PJ, Edvinsson L. The trigeminovascular system and migraine: studies characterizing cerebrovascular and neuropeptide changes seen in humans and cats. *Ann Neurol.* 1993;33(1):48-56.
- Lassen LH, Haderslev PA, Jacobsen VB, Iversen HK, Sperling B, Olesen J. CGRP may play a causative role in migraine. *Cephalalgia.* 2002;22(1):54-61.
- Dodick DW, Goadsby PJ, Silberstein SD, Lipton RB, Olesen J, Ashina M, *et al.* Safety and efficacy of ALD403, an antibody to calcitonin gene-related peptide, for the prevention of frequent episodic migraine: a randomized, double-blind, placebo-controlled, exploratory phase 2 trial. *Lancet Neurol.* 2014;13(11):1100-7.
- Dodick DW, Goadsby PJ, Spierings EL, Scherer JC, Sweeney SP, Grayzel DS. Safety and efficacy of LY2951742, a monoclonal antibody to calcitonin gene-related peptide, for the prevention of migraine: a phase 2, randomized, double-blind, placebo-controlled study. *Lancet Neurol.* 2014;13(9):885-92.
- Bigal ME, Dodick DW, Rapoport AM, Silberstein SD, Ma Y, Yang R, *et al.* Safety, tolerability and efficacy of TEV-48125 for preventive treatment of high-frequency episodic migraine: a multicenter, randomized, double-blind, placebo-controlled, phase 2b study. *Lancet Neurol.* 2015;14(11):1081-90.
- Bigal ME, Edvinsson L, Rapoport AM, Lipton RB, Spierings EL, Diener HC, *et al.* Safety, tolerability and efficacy of TEV-48125 for preventive treatment of chronic migraine: a multicenter, randomized, double-blind, placebo-controlled, phase 2b study. *Lancet Neurol.* 2015;14(11):1091-100.
- Sun H, Dodick DW, Silberstein S, Goadsby PJ, Reuter U, Ashina M, *et al.* Safety and efficacy of AMG 334 for prevention of episodic migraine: a randomised, dou-

- ble-blind, placebo-controlled, phase 2 trial. *Lancet Neurol.* 2016;15(4):382-90.
40. Tepper S, Ashina M, Reuter U, Brandes JL, Dolezil D, Silberstein S, *et al.* Safety and efficacy of erenumab for preventive treatment of chronic migraine: a randomised, double-blind, placebo-controlled phase 2 trial. *Lancet Neurol.* 2017;16(6):425-34.
 41. Deen M, Correnti E, Kamm K, Kelderman T, Papetti L, Rubio-Beltrán E, *et al.* Blocking CGRP in migraine patients – a review of pros and cons. *J Headache Pain.* 2017;18(1):96.
 42. McKeon A, Vincent A. Autoimmune movement disorders. *Handb Clin Neurol.* 2016;133:301-15.
 43. Mohammad SS, Dale RC. Principles and approaches to the treatment of immune-mediated movement disorders. *Eur J Paediatr Neurol.* 2018;22(2):292-300.
 44. Tysnes OB, Storstein A. Epidemiology of Parkinson's disease. *J Neural Transm (Vienna).* 2017;124(8):901-5.
 45. Oertel W, Schultz JB. Current and experimental treatments of Parkinson disease: a guide for neuroscientists. *J Neurochem.* 2016;139(Suppl 1):325-37.
 46. Brundin P, Dave KD, Kordower JH. Therapeutic approaches to target *alpha*-synuclein pathology. *Exp Neurol.* 2017;298(Pt B):225-35.
 47. Graus F, Titulaer MJ, Balu R, Benseler S, Bien CG, Cellucci T, *et al.* A clinical approach to diagnosis of autoimmune encephalitis. *Lancet Neurol.* 2016;15(4):391-404.
 48. Dalmau J, Lancaster E, Martínez-Hernández E, Rosenfeld MR, Balice-Gordon R. Clinical experience and laboratory investigations in patients with anti-NMDAR encephalitis. *Lancet Neurol.* 2011;10(1):63-74.
 49. Suleiman J, Dale RC. The recognition and treatment of autoimmune epilepsy in children. *Dev Med Child Neurol.* 2015;57(5):431-40.
 50. Irani SR, Gelfand JM, Bettecher BM, Singhal NS, Geschwind MD. Effect of rituximab in patients with leucine-rich, glioma-inactivated 1 antibody-associated encephalopathy. *JAMA Neurol.* 2014;71(7):896-900.
 51. Flanagan EP, Drubach DA, Boeve BF. Autoimmune dementia and encephalopathy. *Handb Clin Neurol.* 2016;133:247-67.
 52. Pittock SJ, Palace J. Paraneoplastic and idiopathic autoimmune neurologic disorders: approach to diagnosis and treatment. *Handb Clin Neurol.* 2016;133:165-83.
 53. Pohl D, Alper G, van Haren K, Konberg AJ, Lucchinetti CF, Tenenbaum S, *et al.* Acute disseminated encephalomyelitis: updates on an inflammatory CNS syndrome. *Neurology.* 2016;87(9 Suppl 2):S38-45.
 54. Hebert LE, Weuve J, Scherr PA, Evans DA. Alzheimer disease in the United States (2010-2050) estimated using the 2010 census. *Neurology.* 2013;80(19):1778-83.
 55. Hung SY, Fu WM. Drug candidates in clinical trials for Alzheimer's disease. *J Biomed Sci.* 2017;24(1):47.
 56. Graham WV, Bonito-Oliva A, Sakmar TP. Update on Alzheimer's disease therapy and prevention strategies. *Annu Rev Med.* 2017;68:413-30.
 57. Gilman S, Koller M, Black RS, Jenkins L, Griffith SG, Fox NC, *et al.* Clinical effects of Abeta immunization (AN1792) in patients with AD in an interrupted trial. *Neurology.* 2005;64(9):1553-62.
 58. Farlow MR, Andreasen N, Riviere ME, Vostiar I, Vitaliti A, Sovago J, *et al.* Long-term treatment with active A β immunotherapy with CAD106 in mild Alzheimer's disease. *Alzheimer's Res Ther.* 2015;7(1):23.
 59. Salloway S, Sperling R, Fox NC, Blennow K, Klunk WJ, Raskind M, *et al.* Two phase 3 trials of bapineuzumab in mild-to-moderate Alzheimer's disease. *N Engl J Med.* 2014;370(4):322-33.
 60. Vandenberghe R, Rinne JO, Boada M, Katayama S, Scheltens P, Vellas B, *et al.* Bapineuzumab for mild to moderate Alzheimer's disease in two global, randomized, phase 3 trials. *Alzheimer's Res Ther.* 2016;8(1):18.
 61. Honig LS, Vellas B, Woodward M, Boada M, Bullock R, Borrie M, *et al.* Trial of Solanezumab for mild dementia due to Alzheimer's disease. *N Engl J Med.* 2018;378(4):321-30.
 62. Mo J, Li JY, Yang Z, Liu Z, Feng JS. Efficacy and safety of anti-amyloid- β immunotherapy for Alzheimer's disease: a systematic review and network meta-analysis. *Ann Clin Transl Neurol.* 2017;4(12):931-42.
 63. Bittar A, Sengupta U, Kaye R. Prospects for strain-specific immunotherapy in Alzheimer's disease and tauopathies. *NPJ Vaccines.* 2018;3:9.
 64. Relkin NR, Thomas RG, Rissman RA, Brewer JB, Rafii MS, van Dyck CH, *et al.* A phase 3 trial of IV immunoglobulin for Alzheimer disease. *Neurology.* 2017;88(18):1768-75.
 65. Wilson HJ, Goodfellow JA. GBS100: celebrating a century of progress in Guillain-Barré syndrome. La Jolla, CA: Peripheral Nerve Society; 2016.
 66. Raphaël JC, Chevret S, Hughes RA, Annane D. Plasma exchange for Guillain-Barré syndrome. *Cochrane Database Syst Rev.* 2012;7:CD001798.
 67. Hughes RA, Swan AV, van Doorn PA. Intravenous immunoglobulin for Guillain-Barré syndrome. *Cochrane Database Syst Rev.* 2014;9:CD002063.
 68. Van der Meché FG, Schmidt PI, Dutch Guillain-Barré Study Group. A randomized trial comparing intravenous immune globulin and plasma exchange in Guillain-Barré syndrome. *N Engl J Med.* 1992;326(17):1123-9.
 69. Plasma Exchange/Sandoglobulin Guillain-Barré Syndrome Trial Group. Randomised trial of plasma exchange, intravenous immunoglobulin, and combined treatment in Guillain-Barré syndrome. *Lancet.* 1997;349(9047):225-30.
 70. Wijdicks EFM, Klein CJ. Guillain-Barré syndrome. *Mayo Clin Proc.* 2017;92(3):467-79.
 71. Hu MY, Stathopoulos P, O'Connor KC, Pittock SJ, Nowak RJ. Current and future immunotherapy targets in autoimmune neurology. *Handb Clin Neurol.* 2016;132:511-36.
 72. Benedetti L, Briani C, Franciotta D, Fazio R, Paolasso I, Comi C, *et al.* Rituximab in patients with chronic inflammatory demyelinating polyradiculoneuropathy: a report of 13 cases and review of the literature. *J Neurol Neurosurg Psychiatry.* 2011;82(3):306-8.
 73. D'Amico A, Catteruccia M, De Benedetti F, Vivarelli M, Colucci M, Cascioli S, *et al.* Rituximab in childhood onset idiopathic refractory chronic inflammatory demyelinating polyneuropathy. *Eur J Pediatr Neurol.* 2012;16(3):301-3.
 74. Zhang Z, Zhang ZY, Fauser U, Schluesener HJ. FTY720 ameliorates experimental autoimmune neuritis by inhibition of lymphocyte and monocyte infiltration into peripheral nerves. *Exp Neurol.* 2008;210(2):681-90.
 75. Marsh EA, Hirst CL, Llewelyn JG, Cossburn MD, Reilly MM, Krishnan A, *et al.* Alemtuzumab in the treatment of IVIG-dependent chronic inflammatory demyelinating polyneuropathy. *J Neurol.* 2010;257(6):913-9.
 76. Nobile-Orazio E. Multifocal motor neuropathy. *J Neuroimmunol.* 2001;115(1-2):4-18.
 77. Azulay JP, Blin O, Puget J, Boucraut J, Billé-Turc F, Carles G, *et al.* Intravenous immunoglobulin treatment in patients with motor neuron syndromes associated with anti-GM1 antibodies: a double-blind, placebo-controlled study. *Neurology.* 1994;44(3 Pt 1):429-32.
 78. Van den Berg LH, Kerkhoff H, Oey PL, Franssen H, Mollee I, Vermeulen M, *et al.* Treatment of multifocal motor neuropathy with high dose intravenous immunoglobulins: a double-blind, placebo controlled study. *J Neurol Neurosurg Psychiatry.* 1995;59(3):248-52.
 79. Federico P, Zochodne DW, Hahn AF, Brown WF, Feasby TE. Multifocal motor neuropathy improved by IVIg: randomized, double-blind, placebo-controlled study. *Neurology.* 2000;55(9):1256-62.
 80. Leger JM, Chassande B, Musset L, Meininger V, Bouche P, Baumann N. Intravenous immunoglobulin therapy in multifocal motor neuropathy: a double-blind, placebo-controlled study. *Brain.* 2001;124(Pt 1):145-53.
 81. Van Shaik IN, van den Berg LH, de Haan R, Vermeulen M. Intravenous immunoglobulin treatment for multifocal motor neuropathy. *Cochrane Database Syst Rev.* April 18 (2):CD004429.
 82. Joint task force of the EFNS and the PNS. EFNS/PNS guidelines on management of multifocal motor neuropathy. Report of a joint task force of the European Federation of Neurological Societies and the Peripheral Nerve Society – first revision. *J Peripher Nerv Syst.* 2010;15(4):295-301.
 83. Harbo T, Andersen H, Hess A, Hansen K, Sindrup SH, Jakobsen J. Subcutaneous versus intravenous immunoglobulin in multifocal motor neuropathy: a randomized single-blinded cross-over trial. *Eur J Neurol.* 2009;16(5):631-8.
 84. Harbo T, Andersen H, Jakobsen J. Long-term therapy with high doses of subcutaneous immunoglobulin in multifocal motor neuropathy. *Neurology.* 2010;75(15):1377-80.
 85. Kumar A, Patwa HS, Nowak RJ. Immunoglobulin therapy in the treatment of multifocal motor neuropathy. *J Neurol Sci.* 2017; 375:190-7.
 86. Khalid SI, Ample L, Kely R, Ladha SS, Dardis C. Immune modulation in the treatment of amyotrophic lateral sclerosis: a review of clinical trials. *Front Neurol.* 2017;8:486.
 87. Lee JI, Jander S. Myasthenia gravis: recent advances in immunopathology and therapy. *Expert Rev Neurother.* 2017;17(3):287-99.
 88. Diaz-Manera J, Martínez-Hernández E, Querol L, Klooster R, Rojas-García R, Suárez-Calvet X, *et al.* Long-lasting treatment effect of rituximab in MuSK myasthenia. *Neurology.* 2012;78(3):189-93.
 89. Keung B, Robeson KR, DiCapua DB, Rosen JB, O'Connor KC, Goldstein JM, *et al.* Long-term benefit of rituximab in MuSK autoantibody myasthenia gravis patients. *J Neurol Neurosurg Psychiatry.* 2013;84(12):1407-9.
 90. Iorio R, Damato V, Alboini PE, Evoli A. Efficacy and safety of rituximab for myasthenia gravis: a systematic review and meta-analysis. *J Neurol.* 2015;262(5):1115-9.
 91. Howard JF, Barohn RJ, Cutter GR, Freimer M, Juel VC, Mozaffar T, *et al.* A randomized, double-blind, placebo-controlled phase II study of eculizumab in patients with refractory generalized myasthenia gravis. *Muscle Nerve.* 2013;48(1):76-84.
 92. Howard JF, Utsugisawa K, Benatar M, Murai H, Barohn RJ, Illa I, *et al.* Safety and efficacy of eculizumab in anti-acetylcholine receptor antibody-positive refractory

- generalised myasthenia gravis (REGAIN): a phase 3, randomised, double-blind, placebo-controlled, multicentre study. *Lancet Neurol.* 2017;16(12):976-86.
93. Dhillon S. Eculizumab: a review in generalized myasthenia gravis. *Drugs.* 2018;78(3):367-76.
94. Bernatsky S, Josep L, Pineau CA, Bélisle P, Boivin JF, Banerjee D, *et al.* Estimating the prevalence of polymyositis and dermatomyositis from administrative data: age, sex and regional differences. *Ann Rheum Dis.* 2009;68(7):1192-6.
95. Amato AA, Russell JA. Inflammatory myopathies. In: *Neuromuscular disorders.* NewYork: McGraw Hill; 2008; p. 681.
96. Amato AA, Barohn RJ. Evaluation and treatment of inflammatory neuropathies. *J Neurol Neurosurg Psychiatry.* 2009;80(10):1060-8.
97. Gordon PA, Winer JB, Hoogendijk JE, Choy EH. Immunosuppressant and immunomodulatory treatment for dermatomyositis and polymyositis. *Cochrane Database Syst Rev.* Aug 15;(8):CD003643.
98. Mammen A. Autoimmune muscle disease. *Handb Clin Neurol.* 2016;133:467-84.
99. Dalakas MC, Illa I, Dambrosia JM, Soueidan SA, Stein DP, Otero C, *et al.* A controlled trial of high-dose intravenous immune globulin infusions as treatment of dermatomyositis. *N Engl J Med.* 1993;329(27):1993-2000.