



## PROTOCOL

Bilingual edition English/Spanish

### Semaglutide versus GLP-1 agonists. Effectiveness, safety, and quality of life in patients with diabetes mellitus 2. The SEVERAL study

Semaglutida *versus* agonistas GLP-1. Efectividad, seguridad y calidad de vida en pacientes con diabetes mellitus 2. Estudio SEVERAL

José Seijas-Amigo<sup>1,5,6</sup>, Ángel Salgado-Barreira<sup>2</sup>,  
Rosana Castelo-Domínguez<sup>3</sup>, Mercedes Pereira-Piá<sup>4</sup>,  
Moisés Rodríguez-Mañero<sup>1,5</sup>, José Ramón González-Juanatey<sup>1,5</sup>,  
on behalf of the SEVERAL study group\*

<sup>1</sup>Cardiology Department, Complejo Hospitalario Universitario de Santiago de Compostela, A Coruña. Spain. <sup>2</sup>Unidad de Epidemiología e Investigación Clínica, Universidad de Santiago de Compostela, A Coruña. Spain. <sup>3</sup>Pharmacy Department, Centro de Salud de Ribeira, A Coruña. Spain. <sup>4</sup>Pharmacy Department, Centro de Salud de San Roque, Lugo. Spain. <sup>5</sup>Centro de Investigación Biomédica en Red de Enfermedades Cardiovasculares (CIBERCV). <sup>6</sup>Fundación IDIS de Santiago de Compostela, A Coruña. Spain.

\*ANNEX: The SEVERAL INVESTIGATORS.

## Author of correspondence

José Seijas Amigo  
Hospital Clínico Universitario de Santiago  
Travesía da Choupana s/n.  
15706 A Coruña. Spain.

Email:  
jose.seijas.amigo@sergas.es

Received 7 February 2022;  
Accepted 21 April 2022.  
Early Access date (08/27/2022).  
DOI: 10.7399/fh.13215

## Abstract

**Objective:** The cardiovascular disease is the first cause of deaths in patients with diabetes mellitus 2. The objective is to evaluate and compare the weight loss in patients with diabetes treated with the different GLP-1 receptor agonists for the first time. Secondary endpoints are glycosylated hemoglobin reduction, changes in quality of life and physical activity and the safety of these drugs.

**Method:** It is a postauthorization, multicenter, non-randomized and prospective study. 360 Patients that will start treatment for the first time with GLP-1 receptor agonists will be recruited in 10 centers in the National Health System for a period of 6 months and 44 weeks of follow-up. The primary endpoint will be weight loss achieved with the different GLP-1 receptor agonists and the secondary endpoint will be glycosylated hemo-

## Resumen

**Objetivo:** La enfermedad cardiovascular es la causa principal de muerte en pacientes con diabetes mellitus 2. El objetivo principal es evaluar y comparar prospectivamente la pérdida de peso en pacientes con diabetes mellitus 2 tratados por primera vez con los diferentes análogos de la GLP-1. Como variables secundarias se estudiará reducción de la hemoglobina glicosilada, cambios en calidad de vida y actividad física y la seguridad de estos fármacos.

**Método:** Se trata de un estudio postautorización, multicéntrico, no aleatorizado de seguimiento prospectivo. Se reclutarán 360 pacientes que inicien tratamiento por primera vez con análogos de la GLP1 en 10 centros del sistema público durante un período de 6 meses y un seguimiento de 44 semanas. La variable principal será la pérdida de peso con los

## KEYWORDS

Glucagon Like Peptide 1 receptor; Diabetes Mellitus, Type 2; Weight loss; Obesity; Quality of life; Cardiovascular disease.

## PALABRAS CLAVE

Agonistas de los receptores de glucagón tipo 1 (GLP-1); Diabetes mellitus 2; Pérdida peso; Obesidad; Calidad de vida; Enfermedad cardiovascular.



Los artículos publicados en esta revista se distribuyen con la licencia  
Articles published in this journal are licensed with a  
Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.  
<http://creativecommons.org/licenses/by-nc-sa/4.0/>  
La revista Farmacia no cobra tasas por el envío de trabajos,  
ni tampoco por la publicación de sus artículos.

globin reduction, changes in the quality of life through the EuroQol-5D and changes physical activity through the SF-12 questionnaire, and also the safety of these drugs. The estimate recruitment period will be 6 months, from 1 December 2021 to 1 May 2022. The follow up will finish in December 2022.

**Discussion:** The SEVERAL study will try to provide information about weight loss efficacy, changes in quality of life, physical activity and safety of the GLP-1 receptor agonists in patients with diabetes that start treatment with these drugs in the real life. This study try to compare different GLP-1 receptor agonists in terms of effectiveness and safety for a better posterior election when these drugs are used in patients with diabetes mellitus 2 and obesity.

## Introduction

Diabetes mellitus 2 (DM2) is a metabolic disease characterised by poor glycaemic control caused by increased insulin resistance; thus, there is a need for simple, appropriate, safe, and effective therapies<sup>1</sup>. Furthermore, the leading causes of death in patients with DM2 are cardiovascular (CV) and cerebrovascular complications<sup>1</sup>, which highlights the need for new drugs to reduce the risk of CV events in patients with DM2.

Glucagon-like peptide-1 (GLP-1) is an incretin hormone that is synthesised in intestinal L-cells and secreted in response to meals. It acts by increasing pancreatic insulin secretion in response to glucose, reducing glucagon secretion, and suppressing appetite by acting centrally<sup>2</sup>.

In general, in order to maintain adequate glycaemic control, it is essential to implement lifestyle modifications such as diet and exercise, weight loss, and the use of blood-glucose-lowering drugs. In addition to glycaemic control, several pharmacological therapies, such as GLP-1 agonists (aGLP-1) and renal glucose inhibitors (SGLT-2i), have been shown to provide macrovascular protection, reduce major cardiovascular events (MACE), and reduce hospitalisation due to heart failure (HF)<sup>3,4,5</sup>.

Semaglutide (oral and subcutaneous) is a new generation aGLP-1 that is very similar to human GLP-1<sup>1</sup>. Clinical studies have shown that oral semaglutide is safe, well tolerated, and provides a dose-dependent reduction in HbA1c and body weight according to its use in specific indications<sup>6-10</sup>.

It is difficult to compare the results of different studies on GLP-1 analogues because of differences in design, duration, and variability in the definition of the primary endpoint. Based on the hypothesis that greater weight reductions are achieved with semaglutide than with other types of aGLP-1, this study assesses weight reductions with the different aGLP-1 analogues in patients initiating aGLP-1 treatment in real-life settings. However, due to the high rate of gastrointestinal events and possible treatment abandonment, a further aim is to determine whether weight reduction or possible changes in quality of life and physical activity could lower the rate of treatment failure.

The primary objective of this study is to assess changes in weight loss as measured in kilograms and by percentage, as well as changes in Body Mass Index (BMI) in patients with DM2 treated with semaglutide vs patients treated with other GLP-1 agonists.

Secondary objectives:

- Reductions in HbA1c.
- Changes in quality of life and physical activity (as measured on the EQ-5D-5L and SF-12 questionnaires).
- Safety of these drugs.

## Methods

The SEVERAL study is a multicentre prospective follow-up study with a epidemiological cohort design that includes patients treated with semaglutide (oral and subcutaneous) and patients treated with other GLP-1 analogues (subcutaneous). It is a multidisciplinary study across different levels of health care. The study is promoted and coordinated by a Spanish National Health System (NHS) Cardiology Service. Patients will be recruited from 10 NHS centres. A total of 360 patients will be included in the study (approximately 36 patients per centre). Recruitment will be competitive and will be closed once the total is reached. Based on previous prescription

diferentes análogos de la GLP1 y como variable secundaria se valorarán: reducción de hemoglobina glicosilada, cambios en la calidad de vida y actividad física a través del EuroQol-5D y SF-12 y seguridad. Se ha estimado un período de reclutamiento de 6 meses, desde el 1 de Diciembre 2021 al 1 de Mayo 2022. El seguimiento finalizará en Diciembre de 2022.

**Discusión:** El estudio intentará aportar información sobre la efectividad en pérdida de peso, cambios en calidad de vida, actividad física y seguridad de los análogos de la GLP1 en pacientes con diabetes mellitus 2 que inician tratamiento con estos fármacos en la vida real. Este trabajo pretende comparar los diferentes análogos de la GLP1 en términos de eficacia y seguridad para una posterior mejor elección en la prescripción de estos fármacos en pacientes con diabetes mellitus 2 y obesidad.

statistics, it is estimated that the two groups will be approximately equal (semaglutide is financed for approximately half of the patients who are currently prescribed the drug).

The target population is the DM2 population in the Autonomous Community in which the study will be conducted.

## Selection criteria

### Inclusion criteria

- Being of legal age and more than 18 years of age without an upper limit.
- Being able to understand the study aims and to give consent to participate in the study.
- Prescribed and started treatment with a financed and approved aGLP-1.

### Exclusion criteria

- Diagnosis of any disease, such as proliferative diabetic retinopathy or a family history of thyroid cancer (contraindicated in the Summary of Product Characteristics).
- Gestational diabetes.
- Pregnancy.

### Recruitment phase and selection visit

The approved pharmacists at each centre will be responsible for enrolling patients after they have signed the informed consent form. The assignment of patients to particular therapeutic strategies is not decided in advance by the study protocol, but is determined by standard medical practice. No diagnostic or follow-up intervention will be applied to the patients that is not standard clinical practice.

### Follow-up

Patients will be followed up at the visits listed below, with a window of +/- 2 weeks. Visits 4 and 12 will be by telephone and/or data will be obtained from electronic medical record (when available). If the weight variable is not available for the week 12 visit, patients will be seen in consultations. The week 44 visit will be conducted in the consultation room. Data on the HbA1c variables will be collected according to usual clinical practice and always using the value closest to the consultation (see table 1, Schedule of Visits).

### Data collection

The pharmacists in each centre will be responsible for including patients and collecting data on possible adverse events and different variables (including medication and medical history) through clinical histories and/or telephone interviews. The database will be electronic and anonymised. The study sponsor will be responsible for data validity and custody.

## Statistical analysis

**Sample size.** To achieve the main objective of this study (i.e. to assess weight reduction in patients with DM2 on semaglutide vs patients on other GLP-1 agonists), it is expected that 360 patients will be included during the study period. It has been estimated that approximately 36 patients per

**Table 1.** Schedule of visits

	Inclusion	Visit, week 4 (tel./e-record)	Visit, week 12 (tel./e-record)	Visit, week 44 FINAL
Informed Consent	√			
EQ-5D	√			√
SF-12	√			√
HbA1C/Weight	√		√	√
Safety		√	√	√

e-record: data obtained from electronic medical record; tlf: telephone visit.

centre (10 centres) will be enrolled over a 6-month period. This estimate is based on previous data on the number of monthly approvals at participating centres. Previous studies<sup>11-15</sup> have assessed weight loss in patients with DM2 treated with semaglutide and other GLP-1 agonists, and have described mean weight losses of 3.6 to 4.9 kg in patients on semaglutide vs 0.86 to 2.96 kg in patients on other GLP-1 agonists. Based on these data, and assuming a P-value of 5% as a cutoff for statistical significance, a mean weight loss of 2.5 kg in the GLP-1 agonist group, a mean weight loss of 4.2 kg in the semaglutide group, and a pooled deviation of 3.0 kg, the inclusion of 360 patients will achieve a statistical power of more than 90% to detect differences using the Student *t*-test for independent samples. On the other hand, assuming a possible loss to follow-up of up to 20%, the sample could be reduced by up to 72 patients. In this situation, the statistical power to detect differences in weight loss between the study groups would still be more than 90%. The SPSS 3.0 (Chicago, USA) statistical package was used to estimate statistical power based on the sample size.

Qualitative variables will be expressed as frequency and percentage and quantitative variables will be expressed as mean and standard deviation or as median and interquartile range if they do not conform to normal distribution. Normal distributions will be determined using the Kolmogorov-Smirnov test.

The overall objective will be assessed according to weight loss in each group. Other variables associated with weight loss will be analysed using the Student *t*-test for quantitative variables and the chi-square test for qualitative variables. Multivariate linear regression analyses will be used to adjust for weight loss in relation to the variables shown to be statistically significantly associated in the univariate analysis. Similarly, the specific objectives will be investigated using univariate and multivariate analyses. The coding variables will be categorical, the Student *t*-test or ANOVA will be applied to continuous variables, and the chi-square test will be applied to categorical variables. Multivariate logistic regression will be conducted using significant variables.

#### Questionnaires (see Annexes 1 and 2)

- During the visits indicated in the schedule, the EQ-5D-5L and SF-12 questionnaires will be given out and completed by the participants in consultations.

This study has been approved by the Regional Clinical Research Ethics Committee (CEICm) on December 22, 2021 (No. 2021/471). It has been registered in the Spanish Clinical Trials Register (GESTO) with identifier 0065-2021-OBS (No. AEMPS 21-0022) and it has also been registered at [www.clinicaltrials.gov](http://www.clinicaltrials.gov). The study has been approved by the health authorities.

## Discussion

The SEVERAL study aims to assess potential differences between available GLP-1 agonists using real-life data related to the following aspects: 1) effectiveness in weight reduction; and 2) changes in quality of life, physical activity, and safety and tolerability.

The initial hypothesis is that there are differences between classes regarding weight reduction and tolerability, although an acceptable level

of safety as reported in pivotal studies is to be expected. Such results are believed to be of clinical interest to professionals who prescribe and dispense these promising molecules for the prevention and treatment of patients with diabetes and cardiovascular disease. In addition to semaglutide (oral and subcutaneous), four drugs of the aGLP-1 family are currently available: dulaglutide, exenatide, liraglutide, lixisenatide (subcutaneous). However, no evidence is available on their differences regarding real-life weight reduction and, in particular, to what extent patients adhere to treatment as a result of the high rate of gastrointestinal events outside the setting of clinical trials.

## Clinical trials with GLP-1 analogues

Several clinical trials have been conducted on GLP-1 agonists. All these trials have demonstrated their effectiveness in reducing HbA1c in patients on monotherapy or on combinations with other oral antidiabetic drugs and/or insulin. Dulaglutide was associated with a weight reduction of -0.35 kg to -2.90 kg<sup>8</sup>. In the case of exenatide, 3% of patients experienced at least one period of rapid weight loss (greater than 1.5 kg/wk)<sup>8</sup>. In the case of liraglutide, weight reduction was more significant when the baseline BMI was higher<sup>8</sup>. Lixisenatide was associated with a weight reduction of 1.76 kg to 2.96 kg<sup>8</sup>. Finally, semaglutide was associated with a weight losses of at least 5% and at least 10% in more patients than in those on the active comparators dulaglutide and exenatide, respectively (SUSTAIN 7 and SUSTAIN 3)<sup>12,13</sup>. In the SUSTAIN 6<sup>9,10</sup> trial, weight reduction was -3.6 kg to -4.9 kg. Semaglutide was approved by the FDA for the treatment of obesity in diabetic and non-diabetic patients following the publication of the results of the STEP1 study, which found a weight reduction of -14.9% and a change in baseline weight of -15.3 kg<sup>10</sup>. Subsequently, the STEP 8 trial (weekly subcutaneous semaglutide vs liraglutide) found a weight reduction of -15.8% with semaglutide vs -6.4% with liraglutide<sup>11</sup>.

After assessing these data, it can be concluded that all GLP-1 agonists reduce body weight to varying extents: however, the clinical trials that led to their approval have some limitations that could affect the external validity of the results and their extrapolation to real clinical practice. Each of these trials had different study populations, followed different methodological designs, and differed in relation to the primary endpoint. With the exception of the STEP 1 study<sup>10</sup>, most of these trials were designed to assess HbA1c reduction and cardiovascular events as endpoints, but not to assess weight reductions as endpoints. In contrast to the SEVERAL study, no other study has compared the effectiveness of all GLP-1 agonists.

## Safety of GLP-1 agonists

These weight losses have been accompanied by specific adverse events, the most common being gastrointestinal adverse events, such as nausea, vomiting, diarrhoea, dyspepsia, and constipation<sup>8</sup>. The most recent study on this issue, STEP 8<sup>11</sup>, found that treatment discontinuation with semaglutide was 13.5% and 27.6% with liraglutide. Gastrointestinal events were reported in 84.1% of patients on semaglutide and 82.7% of patients on liraglutide. Most events were mild to moderate, did not lead to permanent discontinuation, were mainly of short duration, and occurred

during dose escalation. These results may indicate a lack of adherence in real life: that is, outside the setting and control to which patients included in clinical trials are exposed. The foregoing suggestion is one of the main motivations for this study (i.e. the assessment of the safety of these drugs in real life).

The SEVERAL study has been designed to collect data on all possible adverse events during the dose escalation and follow-up periods. Data on quality of life and physical activity will also be collected and correlated with potential therapeutic failure and clinical effectiveness.

## Limitations

The present study has several limitations that must be taken into account when interpreting the data. The first limitation is the lack of randomisation, which means that comparative conclusions cannot be drawn: thus, the study merely provides guidance on the current situation, prescription rates, results, and tolerability. If marked differences between classes were to emerge from the present study, a randomised controlled trial could be considered.

Secondly, there is a potentially relevant bias arising from possible loss to follow-up, such that patients with poorer drug tolerability may drop out of the study, thus distorting the results. To avoid this bias, possible loss to follow-up was limited to <20% to calculate sample size. All patients in our Autonomous Community have electronic records, and so the electronic prescription can be checked for medication collection, which will be of great help in validating adherence (as is done in daily clinical practice).

It is worth mentioning that the present working group, which is exclusively made up of professionals from pharmacy units, will minimise inclusion losses as they represent the largest nucleus of dispensing control. Thus, possible bias can be avoided had this study been initiated, for example, by cardiology or endocrinology units, where it would have been more difficult to ensure the correct inclusion of all patients who are prescribed GLP-1 agonists in our Autonomous Community.

In conclusion, this study will attempt to provide information on the effectiveness of GLP-1 agonists on weight reduction and on changes in quality of life, physical activity, and safety in patients with DM2 starting on treatment with these drugs in real life. This study attempts to compare different GLP-1 agonists in terms of effectiveness and safety in order to make better choices in the prescription of these drugs in patients with cardiovascular disease.

## Funding

No funding.

## Acknowledgements

We would like to acknowledge all the main and sub-investigators of the centres participating in the study belonging to the Pharmacy Consultations of SERGAS public centres.

## Conflict of interest

No conflict of interest.

## Bibliography

- Cosentino F, Grant PJ, Aboyans V, Bailey CJ, Ceriello A, Delgado V, et al. Grupo de trabajo de diabetes, prediabetes y enfermedades cardiovasculares de la Sociedad Europea de Cardiología y la European Association for the Study of Diabetes Guía ESC 2019 sobre diabetes, prediabetes y enfermedad cardiovascular, en colaboración con la European Association for the Study of Diabetes (EASD). *Rev. Esp. Cardiol.* 2020;73(5):393-402. DOI: 10.1016/j.recesp.2019.07.013
- Shah N, Abdalla MA, Deshmukh H, Sathyapalan T. Therapeutics for type-2 diabetes mellitus: a glance at the recent inclusions and novel agents under development for use in clinical practice. *Ther Adv Endocrinol Metab.* 2021; 23;12:20420188211042145. DOI: 10.1177/20420188211042145
- Packer M, Anker SD, Butler J, Filippatos G, Pocock SJ, Carson P, et al. Cardiovascular and Renal Outcomes with Empagliflozin in Heart Failure. *N Engl J Med.* 2020;383(15):1413-24. DOI: 10.1056/NEJMoa2022190
- Marso SP, Daniels GH, Brown-Frandsen K, Kristensen P, Mann JF, Nauck MA, et al. Liraglutide and Cardiovascular Outcomes in Type 2 Diabetes. *N Engl J Med.* 2016; 375(4):311-22. DOI: 10.1056/NEJMoa160382
- Gerstein HC, Colhoun HM, Dagenais GR, Diaz R, Lakshmanan M, Pais P, et al. Dulaglutide and cardiovascular outcomes in type 2 diabetes (REWIND): a double-blind, randomised placebo-controlled trial. *Lancet.* 2019;394(10193):121-30. [https://doi.org/10.1016/S0140-6736\(19\)31149-3](https://doi.org/10.1016/S0140-6736(19)31149-3)
- Pratley RE, Aroda VR, Lingvay I, Lüdemann J, Andreassen C, Navarria A, et al. *Lancet Diabetes Endocrinol.* 2018;6(4):275-86. DOI: 10.1016/S2213-8587(18)30024-X
- Davies M, Pieber TR, Hartoft-Nielsen ML, Hansen OKH, Jabbour S, Rosenstock J. Effect of Oral Semaglutide Compared With Placebo and Subcutaneous Semaglutide on Glycemic Control in Patients With Type 2 Diabetes: A Randomized Clinical Trial. *JAMA.* 2017;318(15):1460-70. DOI: 10.1001/jama.2017.14752
- Agencia Española de Medicamentos y Productos Sanitarios (AEMPS). Internet database, CIMA; 2013 [23/02/2022,08/04/2022]. Available at: <https://cima.aemps.es/cima/publico/home.html>
- Marso SP, Holst AG, Vilsbøll T. Semaglutide and Cardiovascular Outcomes in Patients with Type 2 Diabetes. *N Engl J Med.* 2017;376(9):891-2. DOI: 10.1056/NEJMc1615712
- Wilding JPH, Batterham RL, Calanna S, Davies M, Van Gaal LF, Lingvay I, et al. Once-Weekly Semaglutide in Adults with Overweight or Obesity. *N Engl J Med.* 2021;384(11):989. DOI: 10.1056/NEJMoa2032183
- Rubino DM, Greenway FL, Khalid U, O'Neil PM, Rosenstock J, Sørrig R, et al. Effect of Weekly Subcutaneous Semaglutide vs Daily Liraglutide on Body Weight in Adults With Overweight or Obesity Without Diabetes: The STEP 8 Randomized Clinical Trial. *JAMA.* 2022 ;327(2):138-50. DOI: 10.1001/jama.2021.23619
- Pratley RE, Aroda VR, Lingvay I, Lüdemann J, Andreassen C, Navarria A, et al. Semaglutide versus dulaglutide once weekly in patients with type 2 diabetes (SUSTAIN 7): a randomised, open-label, phase 3b trial. *Lancet Diabetes Endocrinol.* 2018;6(4):275-86. DOI: 10.1016/S2213-8587(18)30024-X
- Ahmann AJ, Capehorn M, Charpentier G, Dotta F, Henkel E, Lingvay I, et al. Efficacy and Safety of Once-Weekly Semaglutide Versus Exenatide ER in Subjects With Type 2 Diabetes (SUSTAIN 3): A 56-Week, Open-Label, Randomized Clinical Trial. *Diabetes Care.* 2018;41(2):258-66. DOI: 10.2337/dc17-0417

su estado de salud en el día de hoy.

## Annex the SEVERAL study investigators

José Seijas Amigo<sup>1,5,7</sup>, Ángel Salgado-Barreira<sup>2</sup>, Rosana Castelo-Domínguez<sup>3</sup>, Mercedes Pereira-Piá<sup>4</sup>, Moisés Rodríguez-Mañero<sup>1,5</sup>, Alberto Cordero<sup>5,6</sup>, Begoña Cardeso-Paredes<sup>1,7</sup>, Diego Rodríguez-Penas<sup>1,7</sup>, José Ramón González-Juanatey<sup>1,5</sup>, Agustina Fernández-Pérez<sup>8</sup>, Montserrat Fernández-Montenegro<sup>9</sup>, Ana Belén Ponce-Piñón<sup>10</sup>, Marlen Fernández-Silva<sup>11</sup>, María Teresa Pérez-Álvarez<sup>12</sup>, María del Mar Gago-García<sup>13</sup>, José Manuel Iglesias-Moreno<sup>14</sup>, Marta Rodríguez-Barreiro<sup>15</sup>, María Moure-González<sup>1</sup>, Ana Seoane-Blanco<sup>1</sup>, Rita Soler-Martín<sup>1</sup>, Adrián Paz-Couce<sup>1</sup>

<sup>1</sup>Cardiology Department, Complejo Hospitalario Universitario de Santiago de Compostela, A Coruña. Spain. <sup>2</sup>Unidad de Epidemiología e Investigación Clínica, Universidad de Santiago de Compostela, A Coruña. Spain. <sup>3</sup>Pharmacy Department, Centro de Salud de Ribeira, A Coruña. Spain. <sup>4</sup>Servicio de Farmacia, Centro de Salud de San Roque, Lugo. Spain. <sup>5</sup>Centro de Investigación Biomédica en Red de Enfermedades Cardiovasculares (CIBERCV). <sup>6</sup>Hospital de la Fe de Valencia, Santiago de Compostela, A Coruña. Spain. <sup>7</sup>Fundación IDIS de Santiago de Compostela, A Coruña. Spain. <sup>8</sup>Pharmacy Department, Centro de Salud de Vilalba, Lugo. Spain. <sup>9</sup>Pharmacy Department, Centro de Salud de O Carballiño, Ourense. Spain. <sup>10</sup>Pharmacy Department, Centro de Salud de Fene, A Coruña. Spain. <sup>11</sup>Pharmacy Department, Centro de Salud O Ventorrillo, A Coruña. Spain. <sup>12</sup>Pharmacy Department, Centro de Salud de Culleredo, A Coruña. Spain. <sup>13</sup>Pharmacy Department, Centro de Salud de Ribadeo, Lugo, Spain. <sup>14</sup>Pharmacy Department, Centro de Salud de Valmiñor, Pontevedra. Spain. <sup>15</sup>Pharmacy Department, Centro de Salud Virxe Peregrina, Pontevedra. Spain.

## Annex 1. EQ-5D questionnaire

Please mark with an ☒ X the statement in each section which best describes your state of health today.

### Mobility

- I have no problems walking about ☐
- I have some problems walking about ☐
- I am confined to bed ☐

### Self-Care

- I have no problems with self-care ☐
- I have some problems washing or dressing myself ☐
- I am unable to wash or dress myself ☐

### Everyday Activities (e.g., work, study, housework, family or leisure activities)

- I have no problems performing my everyday activities ☐
- I have some problems performing my everyday activities ☐
- I am unable to perform my everyday activities ☐

### Pain/Discomfort

- I have no pain or discomfort ☐
- I have moderate pain or discomfort ☐
- I have much pain or discomfort ☐

### Anxiety/Depression

- I am not anxious or depressed ☐
- I am moderately anxious or depressed ☐
- I am very anxious or depressed ☐

**Annex 1 (cont.).** EQ-5D questionnaire

To help people describe how good or bad their health is, we have drawn a scale similar to a thermometer on which the best imaginable state of health is marked 100 and 0 for the worst state of health you can imagine.

Please draw a line from the box that says “Your health today,” to the point on the scale that, in your opinion, indicates how good or bad your state of health is today.

**Your  
health  
today**

Best  
imaginable  
health status

100

90

80

70

60

50

40

30

20

10

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

0

Worst  
imaginable  
health status

## Annex 2. SF12 questionnaire

The following questions refer to what you think about your health. Your answers will help us to know how you feel and how well you are able to do your everyday activities.

Please answer each question by ticking a box. If you are unsure how to answer a question, please answer what give the best answer you can.

1. In general, you would say that your health is:

1	2	3	4	5
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Excellent	Very good	Good	Fair	Poor

The following questions refer to activities or things that you might do during a typical day. Does your current health limit you from doing these activities or things? If so, how much?

2. **Moderate exertion**, such as moving a table, hoovering, bowling, or walking for more than 1 hour?

3. Climbing **several** flights of stairs

1	2	3
Yes, limited a lot	Yes, limited a little	No, not limited at all
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

During the **past 4 weeks**, have you had any of the following problems in your work or daily activities **because of your physical health**?

4. **Accomplished less** than you would have liked?

5. Had **to stop doing some tasks** during your work or in your daily activities?

1	2
Yes	No
<input type="checkbox"/>	<input type="checkbox"/>
<input type="checkbox"/>	<input type="checkbox"/>

During the **past 4 weeks**, have you had any of the following problems with your work or in your daily activities **because of an emotional problem** (such as feeling sad, depressed, or nervous)?

6. **Accomplished less** than you would have liked **because of an emotional problem**?

7. Did you do work or activities less **carefully** than usual **due to some emotional problem**?

8. During the **last 4 weeks**, how much did **pain** interfere with your typical work (including work outside the home and housework)?

1	2	3	4	5
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Not at all	A little	Moderately	Quite a lot	Extremely



**Annex 2 (cont.).** SF12 questionnaire

The questions below are about how you have been feeling and how things have been going for you during the **past 4 weeks**. For each question, please give the answer that comes closest to how you have been feeling. During the **past 4 weeks**, for how much of the time...

- |   | 1<br>Always              | 2<br>Almost<br>always    | 3<br>Many<br>always      | 4<br>A few<br>times      | 5<br>Once<br>times       | 6<br>Never               |
|---|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 9. ...have you felt calm and peaceful?        | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 10. ...have you had a lot of energy?          | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 11. ...have you felt down-hearted<br>and sad? | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

12. During the **past 4 weeks**, how much of time have physical health or emotional problems interfered with your social activities (such as visiting friends or family)?

- | 1                        | 2                        | 3                        | 4                        | 5                        |
|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| Always                   | Almost                   | Some<br>always           | Once<br>times            | Never<br>sometimes       |